

# Can corals form aerosol particles through volatile sulphur compound emissions?

Elisabeth Deschaseaux<sup>1</sup>, Graham Jones<sup>1,2</sup>, Branka Miljevic<sup>3</sup>, Zoran Ristovski<sup>3</sup>, Hilton Swan<sup>1</sup>, Petri Vaattovaara<sup>4</sup>

<sup>1</sup> School of Environment, Science and Engineering, Southern Cross University, Lismore NSW 2480 Australia

<sup>2</sup> Marine Ecology Research Centre, Southern Cross University, Lismore NSW 2480 Australia

<sup>3</sup> Queensland University of Technology, Brisbane QLD 4001 Australia

<sup>4</sup> University of Eastern Finland, Kuopio, 70210 Finland

Corresponding author: [e.deschaseaux.10@scu.edu.au](mailto:e.deschaseaux.10@scu.edu.au)

**Abstract.** Acropora dominated coral reefs are a substantial source of atmospheric dimethylsulphide (DMS<sub>a</sub>), one of the most abundant reduced sulphur gases present in the marine boundary layer. DMS is believed to act as a climate regulator of solar radiation and sea surface temperatures through the formation of non-sea-salt sulphate aerosols and cloud condensation nuclei (CCN), although this regulation has not yet been demonstrated. A bubbling chamber experiment was conducted on coral reef seawater containing a branch of *Acropora pulchra*, to investigate whether the coral-generated DMS<sub>a</sub> could be oxidised to non-sea-salt sulphate aerosols under treatment with UV light and O<sub>3</sub>. Results indicated that *A. pulchra* produced significant amounts of dimethylsulphoniopropionate (DMSP) and dissolved DMS although emissions of DMS<sub>a</sub> in the chamber headspace were reduced by the presence of the coral, probably as a result of antioxidant activity in the coral tissue. Significant amounts of carbon disulphide (CS<sub>2</sub>) and ethanethiol (ESH), other sulphur gases that could be involved in CCN formation, were also indicated in the bubbling chamber, most likely from coral production. A decrease in DMS<sub>a</sub> and CS<sub>2</sub> in the presence of UV light and O<sub>3</sub> followed by an occurrence of freshly nucleated nanoparticles (<10nm) suggested that these two sulphur compounds were oxidised and potentially participated in aerosol particle formation and thus could be involved in CCN formation and possibly climate regulation. The study provided insights into the production of sulphur compounds by Acropora dominated coral reefs with potential impact on local climate.

**Key words:** Dimethylsulphide, Coral, Aerosols, Climate.

---

## Introduction

According to the CLAW hypothesis, atmospheric dimethylsulphide (DMS<sub>a</sub>) generated by phytoplankton is oxidised to non-sea-salt (nss) sulphate aerosols which go on to form cloud condensation nuclei (CCN), increasing the albedo of stratocumulus clouds and locally lowering solar radiation and sea surface temperatures (SST) in the ocean (Charlson *et al.*, 1987). Although supported by several studies (Ayers *et al.*, 1991; Modini *et al.*, 2009), the CLAW hypothesis has never been verified. On the other hand, oxidised organic compounds can also participate in newly-formed secondary aerosol particle formation (Vaattovaara *et al.*, 2006).

Research has shown that hard corals, through their symbiotic microalgae, produce DMSP, the main precursor of DMS (Jones *et al.*, 1994; Hill *et al.*, 1995), and emit DMS<sub>a</sub> in chamber experiments on Acropora (Fischer and Jones, 2012). Continuous monitoring of sea surface temperatures (SST) in coral reefs worldwide has shown that pristine reefs within or near the Western Pacific Warm Pool have had fewer reported coral bleaching events relative to reefs in other regions of the world possibly because of an

“ocean thermostat” mechanism that acts to depress warming beyond certain SST thresholds (Kleypas *et al.*, 2008). Research on DMS and aerosol formation suggests that oxidation of DMS<sub>a</sub> from reefs could form nss-sulphate aerosols and thus could contribute to this phenomenon (Jones and Trevena, 2005; Jones *et al.*, 2007; Modini *et al.*, 2009; Jones and Ristovski, 2010).

Bubble bursting from breaking waves in the ocean is the primary source of sea-air exchange and sea spray aerosol production in the atmosphere, constituting a possible source for CCN formation (Modini *et al.*, 2009). Hydroxyl radicals in the atmosphere, which are formed as a consequence of photodissociation of ozone (O<sub>3</sub>) by solar UV, are responsible for the oxidation of gaseous precursors such as DMS into atmospheric aerosols (Andreae and Crutzen, 1997). Thus, bubble bursting, O<sub>3</sub> and UV radiation are three major components in the emission and oxidation of DMS to aerosols.

Between May and June 2011, the CORACE-1 (COral Reef Aerosol Characterization Experiment-1) campaign was conducted on Heron Island coral cay as a collaborative research project between QUT (Queensland University of Technology, Brisbane), SCU (Southern Cross University, Lismore) and UEF (University of Eastern Finland, Kuopio) in order to ascertain whether atmospheric DMS produced by coral reefs could, during bubble bursting and under O<sub>3</sub> and UV radiation, contribute to aerosol particle formation and hence potentially influence local climate (for more background information refer to Swan *et al.* 2012).

## Material and Methods

### Study design and terminology

A bubbling chamber experiment was conducted on seawater collected from the Heron Island Reef flat (23°26'35.80"S/151°54'44.23"E) in which a branch of *Acropora pulchra*, a widely spread coral in the Indo-Pacific and Great Barrier Reef (GBR) (Veron, 2000), was immersed (coral seawater). Three control bubbling chamber experiments were also conducted on coral reef seawater collected at high tide (high tide seawater), low tide (low tide seawater) and from the Heron Island's tap seawater system that pumps water directly from the reef flat (reticulated seawater) (Fig. 1). A control air sample (CAS) was also taken from the "aquaria room" in which the experiment was held. In order to simplify the terminology used for this experiment, "high tide seawater", "low tide seawater" and "reticulated seawater" are referred to as "control seawaters".




Figure 1: Experimental design of the bubbling chamber experiment conducted at Heron Island, May-June 2011, on coral seawater, high tide seawater (HTSW), low tide seawater (LTSW) and reticulated seawater (RSW).

Bubble bursting, simulating sea-air exchange of volatile substances present in the seawater, was intermittently applied to the four types of seawater using dried and filtered compressed air. The air flushed out of the bubbling chamber intermittently went through a particle filter before reaching the reaction chamber in which UV light (40W) and O<sub>3</sub> (200-500ppb) treatments were also applied

intermittently. When filtration was applied, the bubble burst primary particles were removed leaving the gaseous compounds to enter the reaction chamber. When applied, UV and O<sub>3</sub> were simulating and accelerating oxidation processes that may occur to DMS<sub>a</sub> and to other organic compounds that are released from the reefs to the atmosphere during air-sea exchange.

### Sampling

Two types of samples were collected during the experiment: air samples from the headspace of the bubbling chamber simulating atmospheric sulphur emissions from the reef; and water samples from the bubbling chamber, simulating the dissolved sulphur compounds produced within the water column from the reef. Air samples collected onto gold-wool tubes (Kittler *et al.*, 1992) were taken from either the top of the bubbling chamber (before the air was flushed through the reaction chamber) or from the reaction chamber's outlet (Fig.1). Both air and water samples were collected in various conditions: 1) before and after bubbling, 2) with or without air filtration, 3) with or without UV and O<sub>3</sub> treatments and, 4) before and after the coral branch was placed in and taken out of the chamber (coral seawater only). Air samples were collected for atmospheric sulphur compounds. Water samples were collected for total organic and inorganic carbon (TOC and TIC), chlorophyll-a, pheophytin, dissolved DMS (DMS<sub>w</sub>) and DMSP. Temperature, salinity, conductivity and pH were monitored throughout the experiment.

Number and size of primary and secondary particles were determined using a SMPS (scanning mobility particle sizer). Particle chemical properties (composition, volatility, hygroscopicity, oxidised organic fraction) were determined using an Aerodyne ToF-AMS (time-of-flight aerosol mass spectrometer), VH-TDMA (volatilisation and humidification tandem differential mobility analyser) (Fletcher *et al.*, 2007) and UFO-TDMA (ultrafine organic tandem differential mobility analyser) (Vaattovaara *et al.*, 2005) that were placed in-line with the reaction chamber and bubbling chamber.

### Analysis

Sulphur samples were analysed with a purge and trap technique coupled to a gas chromatograph (GC) with a pulsed flame photometric detector (PFPD) using a dual eight-port/six-port two-position manual valve switching system (Swan and Ivey, 1994). Acidified DMSP samples were analysed by alkaline hydrolysis in a purging chamber whereas acidified DMS<sub>w</sub> samples were analysed from the headspace (note: values for

DMSP were corrected for free DMS<sub>w</sub> content). Analysis of TOC and TIC samples was carried out using a total organic carbon analyser. Absorbance of chlorophyll-a and pheophytin samples was determined by spectrophotometry (APHA, 1998).

## Results

Three main peaks (31µM, 39.6µM and 10.8µM) of DMSP and an increasing concentration of DMS<sub>w</sub> (up to 0.3µM) were found in coral seawater containing the branch of *A. pulchra*, whereas DMSP and DMS<sub>w</sub> were not found in control seawaters (Fig. 2).




Figure 2: DMS<sub>w</sub> and DMSP concentrations in coral seawater and control seawaters (RSW, HTSW, LTSW) during the bubbling chamber experiment. Shading shows bubbling periods. CO and CI stand for “coral out” and “coral in” respectively.

DMS<sub>a</sub> as well as two background sulphur compounds, carbon disulphide (CS<sub>2</sub>) and ethanethiol (ESH), were detected in the headspace of the bubbling chamber (Fig. 3). The concentrations of DMS<sub>a</sub>, CS<sub>2</sub> and ESH measured in the “aquaria room” air were much lower than the concentrations of these sulphur compounds contained in the bubbling chamber.




Figure 3: Sulphur gas concentrations in bubbling chamber experiments conducted on coral seawater and control seawaters (RSW, HTSW, LTSW). Atmospheric concentrations of the control air sample (CAS) is also shown. Shading shows bubbling periods. CO and CI stand for “coral out” and “coral in”, respectively.

Although CS<sub>2</sub> (5.2 nmol/m<sup>3</sup>) and ESH (2.8 nmol/m<sup>3</sup>) were detected from the headspace of the chamber prior to applying bubbling, seawater-air exchange of DMS<sub>a</sub> only took place when bubbling was applied. Then 5.5 nmol/m<sup>3</sup> of DMS<sub>a</sub> was measured in the headspace of the bubbling chamber containing the branch of *A. pulchra*. However, five times more DMS<sub>a</sub> and twice as much CS<sub>2</sub> were released from the coral seawater as soon as the coral branch was taken out of the chamber while maintaining constant bubbling.

Generally, the DMS<sub>a</sub> concentration in the headspace of the chamber was similar when purging control seawaters and coral seawater, after the coral branch had been taken out of the chamber. ESH was

present in all experiments but in lower concentration than both DMS<sub>a</sub> and CS<sub>2</sub>. CS<sub>2</sub> was found in all types of seawater except for low tide seawater.

Of particular interest was the finding that DMS<sub>a</sub> and CS<sub>2</sub> tended to decrease as soon as UV and O<sub>3</sub> were applied to the reaction chamber. No decrease in ESH could be recorded as a result of UV and O<sub>3</sub> treatment.

A significant increase in chlorophyll-a and TOC, as well as a decrease in TIC, were found in coral seawater towards the end of the bubbling chamber experiment whereas no variation was observed in control seawaters, and concentrations remained very low (data not shown). Temperature, salinity, conductivity and pH did not vary throughout the entire experiment.

The SMPS data showed that new particle formation occurs when gaseous components released into the chamber headspace upon bubbling were exposed to UV and O<sub>3</sub>. These newly formed particles were too small (count median diameter < 10nm) to be measured by the AMS and therefore their chemical composition has not been determined. The hygroscopicity and volatility profiles of these particles were lower and greater than for sulphates, respectively.

The UFO-TDMA measurements when bubbling, particle filtration, UV and O<sub>3</sub> were applied, showed that the freshly formed secondary particles included at least 50% oxidised organic compounds. The formed ultrafine particles were quickly further oxidised (aged) when they grew bigger in size, due to the production of high level of oxidants into the air.

## Discussion

Bubbling chamber experiments conducted on *Acropora pulchra* and *Acropora* dominated coral reef water provided important information on the production of DMS<sub>a</sub> in coral reef ecosystems and its potential role in aerosol formation and climate regulation.

DMSP was clearly produced by the coral *Acropora pulchra*. The coral-produced DMSP was then rapidly cleaved into DMS<sub>w</sub>, probably as a result of DMSP lyase activity, by either the endosymbiont (Yost and Mitchelmore, 2009) or marine bacteria (Todd *et al.*, 2007) present in the seawater.

The pulsed production of DMSP in coral seawater may mimic coral bleaching events and the loss of symbiotic zooxanthellae (Iglesias-prieto *et al.*, 1992). Eventually, mass release of zooxanthellae in the chamber led to an increase in chlorophyll-a and TOC (data not shown). The decrease in TIC, usually used as a proxy for CO<sub>2</sub>, also suggests an increase in coral bleaching and mortality through a decrease in respiration. This

conclusion was supported by observation of excessive cloudiness of the coral seawater, probably linked to a build up in coral mucus and zooxanthellae in the bubbling chamber. Thus, the present results confirmed previous findings that corals produce significant amounts of DMS and DMSP in coral reef seawater through release of coral mucus containing expelled zooxanthellae (Broadbent and Jones, 2004).

Acropora dominated coral reefs at Heron Island have already been shown to be a significant source of DMS and DMSP (Jones *et al.*, 2007; Fischer and Jones, 2012), and concentrations found in this experiment were similar to what is recorded in the literature. However, the presence of coral-reef-produced CS<sub>2</sub> and ESH was unexpected. Both CS<sub>2</sub> and ESH play an important role in the atmospheric sulphur cycle, with CS<sub>2</sub> oxidising into COS and SO<sub>2</sub>, sulphate compounds that can influence CCN formation as well as global climate and are involved in the formation of acid rain (Yu *et al.*, 2004; Kachina *et al.*, 2006). Both compounds can be produced naturally in the environment (Watts, 2000) but can also be produced as a result of pollution (Yu *et al.*, 2004; Kachina *et al.*, 2006). In this study, bubbling chamber experiments were conducted in a closed chamber and the compressed air that was pumped through the chamber to create bubble bursting was pre-filtered. Also, concentrations of sulphur gases measured from the "aquaria room" were very low relative to that measured in the chamber headspace. Therefore, it is unlikely that CS<sub>2</sub> and ESH came from external pollution and thus, are indicated to be a result of reef production.

However, gas chromatography retention time alone cannot be considered as an absolute means to claim the identity of an analyte and a proper confirmation of identity, using spectral instrumental techniques, needs to be carried out for both CS<sub>2</sub> and ESH.

Meanwhile, emissions of DMS<sub>a</sub>, CS<sub>2</sub> and ESH were similar in coral seawater and control seawaters, suggesting that coral reef seawater was already highly concentrated with these sulphur compounds. However, it appears that the presence of the coral inhibited the release of DMS<sub>a</sub> and CS<sub>2</sub> into the bubbling chamber headspace, supporting the theory that under artificial stressful conditions DMS could be used as an antioxidant within the coral tissue (Sunda *et al.*, 2002; Jones *et al.*, 2007). However, increased concentrations of DMS<sub>a</sub> and CS<sub>2</sub> following the removal of the coral branch could suggest that the bulk of these gases could be entrained within the coral mucus and tissue and were released when the coral was removed from the chamber, constituting an increase in DMS<sub>a</sub> emitted from the water surface.

Bubbling was needed to transfer DMS<sub>w</sub> to the chamber headspace as no DMS<sub>a</sub> was measured from the headspace until bubbling was applied. However, the fact that ESH and CS<sub>2</sub> were detected in the headspace of the chamber prior to apply bubbling

suggests that these two sulphur compounds are more volatile than DMS and thus could be more concentrated in the atmosphere relative to their production rates.

Of particular interest was the decrease of DMS<sub>a</sub> and CS<sub>2</sub> under UV and O<sub>3</sub> treatment which indicated that both sulphur compounds could potentially become oxidised in the atmosphere, participating in secondary aerosol particles and possibly CCN formation (Chin and Davis, 1993; Liss *et al.*, 1997; Moore *et al.*, 2011). In contrast, the fact that ESH remained constant after UV and O<sub>3</sub> exposure suggests that ESH does not contribute to the production of atmospheric aerosols.

It is worth noting that the newly-formed nucleation mode particles included a remarkable and even dominating fraction of oxidised organic compounds. Additionally, hygroscopicity data indicate that newly formed particles might not necessarily originate only from DMS and that other sulphur-containing gaseous precursors or other volatile organic compounds could also be responsible for particle formation and growth. The volatility profile of these particles indicates that sulphates were not the dominant component in these particles.

Further bubbling chamber experiments, as described in this paper, need to be conducted on Acropora corals and artificial seawater to 1) confirm the identity of ESH and CS<sub>2</sub> measured during this first bubbling chamber experiment, 2) see if ESH and CS<sub>2</sub> are a result of coral production and 3) confirm that biogenic DMS<sub>a</sub> and CS<sub>2</sub> are oxidised to aerosol particles and participated in forming CCN. By characterising aerosol particles formed in bubbling chamber experiments such as this one, as well as air masses over coral reefs, we are hoping to eventually ascertain whether reef aerosol emissions can affect local climate.

### Acknowledgement

We would like to thank the Academy of Finland (Kuopio, Finland) and Southern Cross University (Lismore, Australia) including the Marine Ecology Research Centre, the centre for Biogeochemistry and the Division of Research for financial support. Thanks also to the Heron Island Research Station team for their assistance during the CORACE-1 campaign. We are also grateful to the GBR Marine Park Authority for being given permission to collect coral specimens.

### Reference

- Andreae MO, Crutzen PJ (1997) Atmospheric aerosols: Biogeochemical sources and role in atmospheric chemistry. *Science* 276:1052-1058
- APHA (1998) Standard methods for the examination of water and wastewater. Washington, D.C.
- Ayers GP, Ivey JP, Gillett RW (1991) Coherence between seasonal cycles of dimethylsulfide, methanesulfonate and sulfate in marine air. *Nature* 349:404-406

- Broadbent AD, Jones GB (2004) DMS and DMSP in mucus ropes, coral mucus, surface films and sediment pore water from coral reefs in the Great Barrier Reef. *Mar Freshw Res* 55:849-855
- Charlson RJ, Lovelock JE, Andreae MO, Warren SG (1987) Oceanic phytoplankton, atmospheric sulfur, cloud albedo and climate. *Nature* 326:655-661
- Chin M, Davis DD (1993) Global sources and sinks of OCS and CS<sub>2</sub> and their distributions. *Glob Biogeochem Cycles* 7:321-337
- Fischer E, Jones GB (2012) Atmospheric dimethylsulphide production from corals in the Great Barrier Reef and links to solar radiation, climate and coral bleaching. Special Issue of Biogeochemistry (in press)
- Fletcher CA, Johnson GR, Ristovski ZD, Harvey M (2007) Hygroscopic and volatile properties of marine aerosol observed at Cape Grim during the P2P campaign. *Environ Chem* 4:162-171
- Hill RW, Dacey JWH, Krupp DA (1995) Dimethylsulfoniopropionate in reef corals. *Bull Mar Sci* 57:489-494
- Iglesias-prieto R, Matta JL, Robins WA, Trench RK (1992) Photosynthetic response to elevated temperature in the symbiotic dinoflagellate *Symbiodinium microadriaticum* in culture. *Proc. Natl. Acad. Sci. U. S. A.* 89:10302-10305
- Jones G, Curran M, Broadbent A, King S, Fischer E, Jones R (2007) Factors affecting the cycling of dimethylsulfide and dimethylsulfoniopropionate in coral reef waters of the Great Barrier Reef. *Environ Chem* 4:310-322
- Jones G, Ristovski Z (2010) Reef emissions affect climate. *Australasian Science* 26-28
- Jones GB, Curran MAJ, Broadbent AD (1994) Dimethylsulfide in the South Pacific. In: Bellwood, O. C., H. & Saxena, N. (eds.) *Recent advances in marine science and technology '94*. James Cook University of North Queensland Townsville, Queensland 4811 Australia.
- Jones GB, Trevena AJ (2005) The influence of coral reefs on atmospheric dimethylsulfide over the Great Barrier Reef, Coral Sea, Gulf of Papua and Solomon and Bismarck Seas. *Mar Freshw Res* 56:85-93
- Kachina A, Preis S, Kallas J (2006) Catalytic TiO<sub>2</sub> oxidation of ethanethiol for environmentally benign air pollution control of sulphur compounds. *Environ Chem Lett* 4:107-110
- Kittler P, Swan H, Ivey J (1992) An indicating oxidant scrubber for the measurement of atmospheric Dimethylsulphide. *Atm Environ.* 26A:2661-2664
- Kleypas JA, Danabasoglu G, Lough JM (2008) Potential role of the ocean thermostat in determining regional differences in coral reef bleaching events. *Geophys. Res. Lett.* 35:
- Liss PS, Hatton AD, Malin G, Nightingale PD, Turner SM (1997) Marine sulphur emissions. *Philos. Trans R Soc Lond Ser B-Biol. Sci.* 352:159-168
- Modini RL, Ristovski ZD, Johnson GR, He C, Surawski N, Morawska L, Suni T, Kulmala M (2009) New particle formation and growth at a remote, sub-tropical coastal location. *Atmos Chem Phys* 9:7607-7621
- Moore MJK, Furutani H, Roberts GC, Moffet RC, Gilles MK, Palenik B, Prather KA (2011) Effect of organic compounds on cloud condensation nuclei (CCN) activity of sea spray aerosol produced by bubble bursting. *Atm Environ* 45:7462-7469
- Sunda W, Kieber DJ, Kiene RP, Huntsman S (2002) An antioxidant function for DMSP and DMS in marine algae. *Nature* 418:317-320
- Swan HB, Ivey JP (1994) Analysis of atmospheric sulfur gases by capillary gas-chromatography with atomic-emission detection. *HRC-J. High Resolut. Chromatogr.* 17:814-820
- Swan HB, Jones GB, Deschaseaux ESM (2012) Dimethylsulfide, climate and coral reef ecosystems. In: 12th International Coral Reef Symposium, 2012 Cairns.
- Todd JD, Rogers R, Li YG, Wexler M, Bond PL, Sun L, Curson ARJ, Malin G, Steinke M, Johnston AWB (2007) Structural and regulatory genes required to make the gas dimethyl sulfide in bacteria. *Science* 315:666-669
- Vaattovaara P, Huttunen PE, Yoon YJ, Joutsensaari J, Lehtinen KEJ, O'Dowd CD, Laaksonen A (2006) The composition of nucleation and Aitken modes particles during coastal nucleation events: evidence for marine secondary organic contribution. *Atmos Chem Phys* 6:4601-4616
- Vaattovaara P, Rasanen M, Kuhn T, Joutsensaari J, Laaksonen A (2005) A method for detecting the presence of organic fraction in nucleation mode sized particles. *Atmos Chem Phys* 5:3277-3287
- Veron J (2000) Corals of the world. Australian Institute of Marine Science, Townsville.
- Watts SF (2000) The mass budgets of carbonyl sulfide, dimethyl sulfide, carbon disulfide and hydrogen sulfide. *Atm Environ* 34:761-779
- Yost DM, Mitchelmore CL (2009) Dimethylsulfoniopropionate (DMSP) lyase activity in different strains of the symbiotic alga *Symbiodinium microadriaticum*. *Mar Ecol Prog Ser* 386:61-70
- Yu Y, Geyer A, Xie PH, Galle B, Chen LM, Platt U (2004) Observations of carbon disulfide by differential optical absorption spectroscopy in Shanghai. *Geophys Res Lett* 31:L11107